







nanoHUB.org

online simulations and more

The product

$$np = n_i^2 \exp\left(\frac{E_{Fn} - E_{Fp}}{k_B T}\right)$$

suggests that the difference $E_{Fn} - E_{Fp}$ is a measure for the deviation from the equilibrium.

- However, this can not be correct distribution function since it is even in *k*, which means that it suggests that current can never flow in a device.
- The fact that makes it not so unreasonable is that average carrier velocities are usually much smaller than the spread in velocity





nanoHUB.org

• Using this form of the distribution function gives

$$n(\mathbf{r},t) = \frac{1}{V} \sum_{k} f(\mathbf{r},\mathbf{k},t) = N_{C} \exp\left(\frac{E_{Fn} - E_{C0}}{k_{B}T}\right)$$

• In the same manner, one finds that the kinetic energy density per carrier is given by

$$u(\mathbf{r},t) = \frac{1}{2}m * v_d^2 + \frac{3}{2}k_B T$$

The first term on the RHS represents the drift energy due to average drift velocity, and the second term is the well known thermal energy term due to collisions of carriers with phonons



nanoHUB.org online simulations and more

BTE Continued ...

• For this to give the proper average, *f* is normalized as follows:

$$\int d\mathbf{r} \int d\mathbf{v} f(\mathbf{v}, \mathbf{r}, t) = 1$$

• To derive an equation of motion for *f*(**v**,**r**,*t*), it is somewhat easier to consider the particle density

$$n(\mathbf{v},\mathbf{r},t) = Nf(\mathbf{v},r,t)$$

where

$N = \int d\mathbf{r} \int d\mathbf{v} n(\mathbf{v}, \mathbf{r}, t) = Total \# of particles$

 The density n(v,r,t) should satisfy a continuity equation in the 6D phase space defined by

 $x, y, z, v_x, v_y, v_z \rightarrow$ Independent variables





BTE Continued ...
BTE Continued ...
which is written more compactly as:

$$\nabla \cdot \mathbf{j} = \mathbf{v} \cdot \nabla_r n + \frac{\mathbf{F}}{m} \nabla_v n$$
• Particle balance is therefore:

$$\int_{V} d\mathbf{r} d\mathbf{v} \left(\frac{\partial n}{\partial t} + \mathbf{v} \cdot \nabla_r n + \frac{\mathbf{F}}{m} \cdot \nabla_v n - \frac{\partial n}{\partial t} \Big|_{Coll} - \frac{\partial n}{\partial t} \Big|_{G-R} \right) = 0$$
Normalizing, we get the classical form of the Boltzmann transport equation:

$$\frac{\partial f(\mathbf{r}, \mathbf{v}, t)}{\partial t} = -\mathbf{v} \cdot \nabla_r f - \frac{\mathbf{F}}{m} \cdot \nabla_v f + \frac{\partial f}{\partial t} \Big|_{Coll} + \frac{\partial f}{\partial t} \Big|_{G-R}$$
First two terms on the rhs are the streaming terms







nanoHUB.org
online simulations and moreCollision Integral Cont'd ...(C) Boltzmann Equation with Collision IntegralThe sum over final states k' may be converted to an integral
due to the small volume of k-space associated with each
state: $\sum_{k'} \rightarrow \frac{V}{8\pi^3} \int dk'$ The BTE becomes: $\frac{\partial f_k}{\partial t} + \frac{1}{\hbar} \nabla_k E \cdot \nabla_r f_k + \frac{F}{\hbar} \nabla_k f_k =$
 $\frac{V}{8\pi^3} \int dk \{f_k [1 - f_k] \Gamma_{k'k} - f_k [1 - f_{k'}] \Gamma_{kk'}\}$



nanoHUB.org online simulations and more

Fermi Golden Rule

Additional Notes: Time dependent perturbation theory

• Assume the Hamiltonian may be decomposed as $H=H_0+V_s$, where H_0 is the Hamiltonian of the perfect crystal (described by Bloch states), $V_s(\mathbf{r},t)$ is a small random potential. If $V_s << H_0$, then it is a good approximation to expand the solution (with random part) in terms of unperturbed eigenstates:

$$H_0 \Psi_k = E_k \Psi_k; \quad \Psi_k^0(\mathbf{r}, t) = \Psi_k(\mathbf{r}) e^{-iE_k t/\hbar}$$

• Expand actual solution in terms of these orthonormal functions:

$$\Psi(\mathbf{r},t) = \sum_{k} c_{k}(t) \Psi_{k}(\mathbf{r}) e^{-iE_{k}t/\hbar}$$



nanoHUB.org online simulations and more	Fermi Golden Rule Cont'd
H_0 part cancels with phase factor on RHS	
$V_{s\sum_{k}} c_{k}(t) \psi_{k}(\mathbf{r}) e^{-iE_{k}t/\hbar} = i\hbar \sum_{k} \frac{\partial c_{k}(t)}{\partial t} \psi_{k}(\mathbf{r}) e^{-iE_{k}t/\hbar}$	
• Multiply both sides by $\psi_{k_0'}({ m r}) e^{-i E_{\kappa_0} t/\hbar}$ and integrate	
$i\hbar \frac{\partial \boldsymbol{c}_{k_0'}(t)}{\partial t} = \sum_k \boldsymbol{c}_k$	$(t)\langle {m k}_0' ig {m V}_{ m s} ig {m k} angle {m e}^{-i ({m E}_{{m k}_0} - {m E}_{m k})t/\hbar}$
where the <i>matrix elemen</i> as	t, using Dirac notation, is defined
$\langle \mathbf{k}_0' \mathbf{V}_s \mathbf{k} \rangle = \int d\mathbf{r} \psi_{\mathbf{k}_0'}^* \mathbf{V}_s(\mathbf{r}, t) \psi_{\mathbf{k}'}$	
	Network for Computational Nanotechnology

Fermi Golden Rule Cont'd ...

• Assume sufficiently weak scattering that $c_{ko} \approx 1$, and $c_{k \neq ko} \approx 0$ for all time. The dominant term in the sum is:

$$i\hbar \frac{\partial c_{k_0'}(t)}{\partial t} = c_{k_0}(t) \langle k_0' | V_s | k_0 \rangle e^{-i(E_{k_0} - E_{k_0})t/\hbar}$$

which integrates to

nanoHUB.org

online simulations and more

$$c_{K_{0}}(t) = \frac{1}{i\hbar} \int_{0}^{t} dt' \langle K_{0}' | V_{s} | k_{0} \rangle e^{-i(E_{k_{0}} - E_{k_{0}})t'/\hbar} + c_{K_{0}'}(0)$$

• Suppose V(r,t) may be Fourier decomposed, so that

$$V_{s}(\mathbf{r},t) = V_{s}(\mathbf{r})e^{\pm i\alpha}$$

Note that this form of $V(\mathbf{r},t)$ may correspond to interaction with lattice vibrations or with optical excitation.



nanoHUB.org

line simulations and more

• Substituting for *c* and taking the magnitude squared gives

$$P_{k_0 K_0} = \lim_{t \longrightarrow \infty} \frac{1}{\hbar^2} |V_s^{k_0 K_0}|^2 \left(\frac{\sin(\Lambda t)}{\Lambda t}\right)^2 t^2$$

where asymptotically

$$\lim_{t \longrightarrow \infty} \left(\frac{\sin(\Lambda t)}{\Lambda t} \right)^2 = 2\pi \delta(\Lambda) / t = 2\pi \hbar \delta(E_{k_0} - E_{k_0} \mp \hbar \omega) / t$$

This gives the famous **Fermi's Golden Rule** (droping 0's index)

$$\Gamma_{kk'} = \frac{P_{kk'}}{t} = \frac{2\pi}{\hbar} \left| V_s^{kk'} \right|^2 \delta(E_{k'} - E_k \mp \hbar \omega)$$

• Assumptions made:

(1) Long time between scattering (no multiple scattering events)

(2) Neglect contribution of other c's (Collisional broadening ignored)

NCN